


Mathcad Selections from Numerical Recipes Function Pack

Platform: Windows

Requires Mathcad PLUS 5.0 or higher, 4 MB hard disk space

Available for immediate download (size 2293261 bytes) or ground shipment




This on-line Function Pack contains over 140 useful functions from the second edition of Numerical Recipes, the best-selling book by William H. Press, William T. Vetterling, Saul A. Teukolsky, and Brian P. Flannery. Once installed, these functions become part of the Mathcad PLUS function set, further enhancing the power of Mathcad. A Mathcad Electronic Book is included to document the functions and provide background information and working examples. You'll benefit from easier, quicker access to ideas and numerical algorithms -- all without the need for programming or compiling code.

[Table of Contents](#)

[Product Sample](#)

[Back to Product List](#)




Use the added functionality in this function pack to simulate annealing methods. The anneal function is used in solving the traveling salesman problem.

Topics include: Two-dimensional Polynomial and Spline Interpolation, Laguerre's Method of Finding Roots, Power Spectra, Wavelet Transforms, Chebyshev and Rational Function Approximations, Monte Carlo Integration, and much more. Applications include: Circuit Analysis, Predator-Prey Interaction, and Prediction of Stock Market Averages.

Mathcad Selections from Numerical Recipes Function Pack

TABLE OF CONTENTS (page 1 of 6)



Solutions of Linear Algebraic Equations

- Gauss-Jordan Elimination
- LU Decomposition and its Applications
- Tridiagonal Systems of Equations
- Iterative Improvement of a Solution to Linear Equations
- Singular Value Decomposition

Interpolation and Extrapolation

- Polynomial Interpolation and Extrapolation
- Rational Function Interpolation and Extrapolation
- Coefficients of the Interpolating Polynomial
- Interpolation in Two or More Dimensions

Integration of Functions

- Improper Integrals
- Gaussian Quadratures and Orthogonal Polynomials
- Multidimensional Integrals

Evaluation of Functions


- Polynomials and Rational Functions
- Chebyshev Approximation
- Polynomial Approximation from Chebyshev Coefficients
- Pade Approximants
- Rational Chebyshev Approximation

[Product Sample](#)

[Back to Product List](#)

Mathcad Selections from Numerical Recipes Function Pack

TABLE OF CONTENTS (page 2 of 6)



Special Functions

The Logarithm of the Gamma Function, Binomial Coefficients, Beta Function, Factorials
Incomplete Gamma Functions, Complementary Error Function
Exponential Integrals
Incomplete Beta Function
Bessel Functions of Fractional Order
Spherical Harmonics
Fresnel Integrals, Cosine and Sine Integrals
Elliptic Integrals and Jacobian Elliptic Functions
Hypergeometric Functions

Random Numbers

Exponential and Normal Deviates
Gamma, Poisson, Binomial Deviates
Generation of Random Bits
Quasi-Random Sequences
Adaptive Monte Carlo

Root Finding and Nonlinear Sets of Equations

Bisection Method
Ridder's Method
Brent's Method
Roots of Polynomials
Globally Convergent Methods for Nonlinear Systems of Equations

Minimization or Maximization of Functions


Parabolic Interpolation and Brent's Method in One Dimension
Downhill Simplex Method in Multidimensions
Direction Set Methods in Multidimensions
Conjugate Gradient Methods in Multidimensions
Variable Metric Methods in Multidimensions
Linear Programming and the Simplex Method
Simulated Annealing Methods

[Product Sample](#)

[Back to Product List](#)

Mathcad Selections from Numerical Recipes Function Pack

TABLE OF CONTENTS (page 3 of 6)



Eigensystems

Jacobi Transformations of a Symmetric Matrix
Eigenvalues and Eigenvectors of a Tridiagonal Matrix

Fast Fourier Transform

Cosine Transform
FT in Two or More Dimensions

Fourier and Spectral Applications

Linear Prediction and Linear Predictive Coding
Power Spectrum Estimation
Spectral Analysis of Unevenly Sampled Data
Wavelet Transforms

Statistical Description of Data

Moments of a Distribution
Do Two Distributions have the Same Means or Variances?
Are Two Distributions Different?
Contingency Table Analysis of Two Distributions
Linear Correlation
Nonparametric or Rank Correlation
Do Two-Dimensional Distributions Differ?

Modeling of Data


Fitting Data to a Straight Line
Straight-Line Data with Errors in Both Coordinates
General Linear Least Squares
Nonlinear Models
Robust Estimation

[Product Sample](#)

[Back to Product List](#)

Mathcad Selections from Numerical Recipes Function Pack

TABLE OF CONTENTS (page 4 of 6)



Integration of Ordinary Differential Equations

Runge-Kutta Method
Richardson Extrapolation and the Bulirsch-Stoer Method
Stiff Sets of Equations

Boundary Value Problems, Integral Equations, Partial Differential Equations

The Shooting Method
Fredholm Equations of the Second Kind
Volterra Equations
Relaxation Methods for Boundary Value Problems

Index of Applications

accuracy of polynomial interpolation
adaptive Monte Carlo integration
benefits of job training
bidding for a contract
center of mass of a inhomogeneous cylinder
center of mass of a weighted board
circuit analysis
convolution of functions
critically damped spring action
displacement of a beam
estimating the frequency of an unknown signal
estimating unknown values in a surface or contour plot

[Product Sample](#)

[Back to Product List](#)

Mathcad Selections from Numerical Recipes Function Pack

TABLE OF CONTENTS (page 5 of 6)

evaluating nasty integrals

exact polynomial derivatives

finding space in a parking garage

finding the balance in a savings account

fitting lines of least absolute deviation

grading on a curve

heat expansion in an unknown metal

heat in a semi-infinite plate

hypergeometric function identities

improving a solution

interpolating to find local minima

intersection points of hypersurfaces

intersections of planetary orbits

kernels of linear transformations

least-squares solution to a differential equation

males vs. females on the SAT


maps and symmetric matrices

maximizing profit on a housing project

motion of a driven spring

optimal location for a new restaurant

pendulum motion



[Product Sample](#)

[Back to Product List](#)

Mathcad Selections from Numerical Recipes Function Pack

TABLE OF CONTENTS (page 6 of 6)

Poisson equation on a square

polynomial approximations to nonpolynomial functions

predator-prey interaction

predicting stock market averages

projectile motion

quality control in manufacturing

sending coded messages

series solutions to differential equations

solution to Legendre's differential equation

sub-random sequences

testing the effectiveness of medication

the distribution of galaxies


the traveling salesman problem

time to failure of a system of components

timing differences with sparse matrices

transforming a 2-dimensional image

vibrations of a circular membrane




[Product Sample](#)

[Back to Product List](#)

Mathcad Selections from Numerical Recipes Function Pack


SAMPLE PAGE (page 1 of 3)



Power Spectrum Estimation

We give an example where we compute the power spectrum estimation of a signal consisting of two sine waves with different frequencies, plus some white noise, which we create using **gasdev** (another function in the Function Pack).

```
N := 29 - 1  
Δ := .1  
f2 := .53  
datai := sin(2·π·ti·f1) + sin(2·π·ti·f2)  
i := 0..N  
ti := Δ·i  
f1 := .57  
data := data + gasdev(N + 1)
```



We compute the power spectrum with 20, 85 and 250 coefficients.

```
M := 1024  
j := 0..M  
xj :=  $\frac{j}{M \cdot 2}$   
E20 := evlmem(x, data, 20)  
E85 := evlmem(x, data, 85)  
E250 := evlmem(x, data, 250)
```




[Table of Contents](#)

[Back to Product List](#)

Mathcad Selections from Numerical Recipes Function Pack

SAMPLE PAGE (page 2 of 3)

To see the interesting range of frequencies clearly, we graph only the range of frequencies containing **f1** and **f2**.





[Table of Contents](#)

[Back to Product List](#)

Mathcad Selections from Numerical Recipes Function Pack

SAMPLE PAGE (page 3 of 3)



We can see that with only 20 coefficients we cannot distinguish between the two peaks corresponding to the different frequencies. With 85 coefficients the peaks start to separate, and with 250 coefficients they are quite distinct, although the background noise is starting to show false peaks, visible with the logarithmic scale.

[Table of Contents](#)

[Back to Product List](#)